arXiv:math/0505573v1 [math.GR] 26 May 2005

THOMPSON'S GROUP F IS NOT KÄHLER

TERRENCE NAPIER* AND MOHAN RAMACHANDRAN

Dedicated to Ross Geoghegan in honor of his 60th birthday.

Abstract. The purpose of this note is to prove that Richard Thompson's group F and variants of it studied by Ken Brown are not Kähler groups.

Introduction

The purpose of this note is to prove the following:

Theorem 0.1. *Thompson's group F and the generalizations $F_{n,\infty}$ and F_n for $n = 2, 3, 4, \ldots$ are not Kähler.*

Theorem 0.1 answers a question of Ross Geoghegan (see Ken Brown's paper in these proceedings [Br2]).

Thompson discovered the group F in 1965 in the context of his work in mathematical logic. Brown and Geoghegan [BrGe] determined that F is of type FP_∞, thus making F the first known example of a torsion free group of type FP_∞ which is not of type FP. The group F has also appeared in homotopy theory [FH]. For more details on the properties and history of Thompson's groups F, $F_{n,\infty}$, and F_n, the reader may refer to [CFP], [Br1], [BG], and [BrGe].

A finitely presented group is called a *Kähler group* if it is the fundamental group of a compact Kähler manifold. A central problem in the study of the topology of compact Kähler manifolds (for example, smooth projective varieties) is that of determining which groups are Kähler groups. For example, according to Hodge theory, the Abelianization of a Kähler group must be of even rank (see, for example, [W]). By [ArBR], a Kähler group has at most one end. If M is any Hopf surface, then M is a non-Kähler compact complex manifold which has fundamental group $\mathbb{Z}$ and 2-ended universal covering $\mathbb{C}^2 \setminus \{0\}$. The Heisenberg group H of 3×3 upper triangular integer matrices with diagonal entries 1

Date: May 25, 2005.

2000 *Mathematics Subject Classification.* 20F65, 32J27.

Key words and phrases. HNN extension, Riemann surface.

*Research partially supported by NSF grant DMS0306441.

has exactly one end and its Abelianization is of rank 2, but H is not a Kähler group (see, for example, [JR] or [Ar]). On the positive side, any finite group is the fundamental group of a smooth projective variety. For any positive integer g, the group $\langle a_1, a_2, \ldots, a_{2g} \mid [a_1, a_{g+1}] \cdot \cdots \cdot [a_g, a_{2g}] = 1 \rangle$ is the fundamental group of a curve of genus g. In fact, much more subtle examples are now known to exist; for example, Toledo's examples of Kähler groups which are not residually finite [T]. The problem is, of course, related to that of determining which groups are fundamental groups of smooth projective varieties. In fact, there are no known examples of Kähler groups which are not also fundamental groups of smooth projective varieties. For more details on the study of Kähler groups, the reader may refer to the surveys [AmBCKT] and [Ar].

We give two proofs that Thompson's group F is not Kähler. The first proof (Section 1) relies on the particular properties of F. The second proof (Section 2) gives the following more general fact which may eventually yield a proof that other groups of interest are not Kähler:

Theorem 0.2. *Let G be a group which satisfies the following:*

- (i) *Any nontrivial normal subgroup contains the commutator subgroup G_{com} (i.e. every proper quotient of G is Abelian), and*
- (ii) *G is a properly ascending HNN extension.*

Then G and any group containing G as a subgroup of finite index are not Kähler.

Since $F_{n,\infty}$ ($F = F_{2,\infty} = F_2$) satisfies the above conditions (i) and (ii) and $F_{n,\infty}$ is a (normal) subgroup of finite index in F_n for every $n \geq 2$ (see, for example, [CFP] and [BG]), Theorem 0.1 is a consequence of Theorem 0.2.

Acknowledgement. We would like to thank Ken Brown and Ross Geoghegan for answering our questions about Thompson's group F and Ross Geoghegan for asking the question that this paper answers. We would also like to thank John Meier for trying to teach us right from wrong in geometric group theory. Finally, we would like to thank the referee for helpful comments.

1. First proof

For the purposes of this note, X will denote a connected compact Kähler manifold and C will denote a connected compact curve (i.e. a compact 1-dimensional complex manifold). A remark we will use without comment is a nonconstant holomorphic map from X to C is surjective and open.

Let $f : X \to C$ be a surjective holomorphic map with connected fibers. Let $\{ p_1, \ldots, p_r\}$ be the set of critical values of f and let m_i be the greatest common divisor of the multiplicities of the components of the divisor $f^{-1}(p_i)$ for each $i = 1, \ldots, r$. Let C^{orb} be the 2-orbifold with underlying topological space C and singular points $p_1, \ldots, p_r$ of order $m_1, \ldots, m_r$, respectively. Note that C^{orb} has a unique structure of a complex 1-orbifold so that the map f is holomorphic. We recall that, in this situation (see [Ct], [Si]), the *orbifold fundamental group* $\pi_1^{\mathrm{orb}}(C)$ is the quotient of $\pi_1(C - \{ p_1, \ldots, p_r \})$ by the normal closure of $\{ \gamma_i^{m_i} \mid i = 1, \ldots, r \}$, where γ_i is a simple loop around the point p_i for each i.

Let $G = \pi_1(X)$. The kernel K of the surjective homomorphism $\pi_1(X) \to \pi_1^{\mathrm{orb}}(C)$ is the image in G of the fundamental group of a general fiber of f. In particular, K is a finitely generated normal subgroup of G. If the underlying topological surface of C has positive genus, then $\pi_1^{\mathrm{orb}}(C)$ is $\mathbb{Z}^2$ or a cocompact Fuchsian group. Summing up, we have the following:

Lemma 1.1. *For $G = \pi_1(X)$, we have an exact sequence*

$$1 \to K \to G \to \pi_1^{orb}(C) \to 1,$$

where K is finitely generated. If C has positive genus, then $\pi_1^{orb}(C)$ is either $\mathbb{Z}^2$ or a cocompact Fuchsian group.

In both proofs of Theorem 0.1, the main point leading to a contradiction is that, if $F = \pi_1(X)$ for X Kähler, then X admits such a mapping $f : X \to C$. In the first proof, the main features of the group F which we will use are (see [CFP]):

1. F is torsion free;
2. The commutator subgroup F_{com} is *not* finitely generated;
3. Any nontrivial normal subgroup contains F_{com}; and
4. The Abelianization F/F_{com} is $\mathbb{Z}^2$.

First proof of Theorem 0.1. If $F = \pi_1(X)$ for some connected compact Kähler manifold X, then, since the Abelianization F/F_{com} of F is $\mathbb{Z}^2$ (property 4), the Albanese variety is an elliptic curve E. Stein factoring the Albanese map $h : X \to E$, we get a commutative diagram of surjective holomorphic maps

$$\begin{tikzcd} X \arrow[r, "h"] \arrow[dr, "f"'] & E \\ & C \arrow[u, "g"'] \end{tikzcd}$$

where C is a compact curve, f has connected fibers, and g has finite fibers. In particular, C is of positive genus. Applying Lemma 1.1, we get the exact sequence

$$1 \to K \to F = \pi_1(X) \to \pi_1^{\mathrm{orb}}(C) \to 1,$$

where K is finitely generated. Thus $F/K \cong \pi_1^{\mathrm{orb}}(C)$. Now F is not cocompact Fuchsian. For, if F were cocompact Fuchsian, then, since F is torsion-free (property 1), F would be the fundamental group of a curve of genus at least 2 and hence F would have Abelianization of rank at least 4 (contradicting property 4). F also cannot be isomorphic to $\mathbb{Z}^2$ because the subgroup F_{com} is not finitely generated (property 2). Thus K is nontrivial and, therefore, K contains the commutator subgroup F_{com} (property 3). Therefore $\pi_1^{\mathrm{orb}}(C)$ is a quotient of the Abelianization $F/F_{\mathrm{com}} \cong \mathbb{Z}^2$. In particular, since cocompact Fuchsian groups are non-Abelian, $\pi_1^{\mathrm{orb}}(C)$ must be isomorphic to $\mathbb{Z}^2$ and hence cannot be isomorphic to a *proper* quotient of $\mathbb{Z}^2$. Thus we must have $\pi_1^{\mathrm{orb}}(C) = F/F_{\mathrm{com}}$; that is, $K = F_{\mathrm{com}}$. But this is impossible because F_{com} is not finitely generated (property 2). □

2. Second proof

We first consider the following:

Theorem 2.1. *Given a connected compact Kähler manifold X and an exact sequence*

$$(*) \qquad 1 \to N \to \pi_1(X) \xrightarrow{\rho} \mathbb{Z} \to 1$$

with N not finitely generated, we get an exact sequence

$$(**) \qquad 1 \to K \to \pi_1(X) \to \Gamma \to 1,$$

where Γ is a cocompact Fuchsian group of a curve of positive genus and K is finitely generated.

Proof. By Theorem 4.3 of [NR], there is a surjective holomorphic map with connected fibers to a curve of positive genus $f : X \to C$ and a factorization

$$\begin{tikzcd} \pi_1(X) \arrow[r, "f_*"] \arrow[dr, "\rho"'] & \pi_1^{\mathrm{orb}}(C) \arrow[d, "\rho'"] \\ & \mathbb{Z} \end{tikzcd}$$

By Lemma 1.1, we have an exact sequence $1 \to K \to \pi_1(X) \to \pi_1^{\mathrm{orb}}(C) \to 1$, where K is finitely generated. Since K is finitely generated but N is not, $\ker(\rho') \cong N/K$ is not finitely generated. It follows that $\Gamma = \pi_1^{\mathrm{orb}}(C)$ is a cocompact Fuchsian group, since every subgroup of $\mathbb{Z}^2$ is finitely generated. ☐

Proof of Theorem 0.2. It suffices to prove that G is not Kähler, because any finite covering of a compact Kähler manifold is compact Kähler. If there exists a connected compact Kähler manifold X with $\pi_1(X) = G$, then, since G is a properly ascending HNN extension (property (ii)), we get an exact sequence of the form $(*)$ as in Theorem 2.1 and, therefore, we get an exact sequence of the form $(**)$. If K is nontrivial, then K contains the commutator subgroup (property (i)). But a cocompact Fuchsian group cannot be Abelian, so we arrive at a contradiction. If K is trivial, then G is cocompact Fuchsian and, therefore, by Malcev's theorem, residually finite; i.e. for any element $g \neq 1$, there is a finite homomorphic image of G in which the image of g is not the identity (for an elementary proof, see [Al]). Taking $g \in G_{\mathrm{com}} \setminus \{1\}$, we get a non-Abelian finite homomorphic image. Thus we again arrive at a contradiction to property (i). Therefore, G is not Kähler. ☐

References

[Al] R. Alperin, *An elementary account of Selberg's lemma*, L'Enseignement Math. **33** (1987), 269–273.

[AmBCKT] J. Amorós, M. Burger, K. Corlette, D. Kotschick, D. Toledo, *Fundamental Groups of Compact Kähler Manifolds*, Math. Surveys and Monographs, Vol. 44, American Mathematical Society, Providence, 1996.

[Ar] D. Arapura, *Fundamental groups of smooth projective varieties*, Current Topics in Complex Algebraic Geometry (Berkeley, CA, 1992/93), 1–16, Math. Sci. Res. Inst. Publ., 28, Cambridge Univ. Press, Cambridge, 1995.

[ArBR] D. Arapura, P. Bressler, M. Ramachandran, *On the fundamental group of a compact Kähler manifold*, Duke Math. Jour **68** (1992), 477–488.

[BG] M. Brin, F. Guzmán, *Automorphisms of generalized Thompson groups*, J. of Algebra **203** (1998), 285–348.

[Br1] K. S. Brown, *Finiteness properties of groups*, J. Pure Appl. Algebra **44** (1987), 45–75.

[Br2] K. S. Brown, *The homology of Richard Thompson's group F*, Proceedings of the Conferences on Probabilistic and Asymptotic Aspects of Group Theory and Topological Aspects of Group Theory (Nashville, 2004), Contemporary Mathematics, American Mathematical Society, Providence.

[BrGe] K. S. Brown, R. Geoghegan, *An infinite-dimensional torsion-free FP_∞ group*, Invent. Math. **77** (1984), no. 2, 367–381.

[CFP] J. W. Cannon, W. J. Floyd, W. R. Parry, *Introductory notes on Richard Thompson's groups*, Enseign. Math. **42** (1996), no. 3-4, 215–256.

[Ct] F. Catanese, *Fibred Kähler and quasi-projective groups*, Special issue dedicated to Adriano Barlotti, Adv. Geom. 2003, suppl., S13–S27.

[FH] P. Freyd, A. Heller, *Splitting homotopy idempotents, II*, J. Pure Appl. Algebra **89** (1993), 93–106.

[JR] F. Johnson, E. Rees, *On the fundamental group of a complex algebraic manifold*, Bull. London Math. Soc. **19**, no. 5, (1987), 463–466.

[NR] T. Napier, M. Ramachandran, *Hyperbolic Kähler manifolds and proper holomorphic mappings to Riemann surfaces*, Geom. Funct. Anal. **11** (2001), 382–406.

[Si] C. Simpson, *The ubiquity of variations of Hodge structure*, Complex geometry and Lie theory (Sundance, 1989), 329–348, Proc. Sympos. Pure Math., Vol. 53, Amer. Math. Soc., Providence, RI, 1991.

[T] D. Toledo, *Projective varieties with nonresidually finite fundamental group*, Publ. Math. IHES **77** (1993), 103–119.

[W] R. O. Wells, *Differential Analysis on Complex Manifolds*, Grad. Texts in Math., Vol. 65, Springer, Berlin-Heidelberg-New York, 1980.

Department of Mathematics, Lehigh University, Bethlehem, PA 18015
E-mail address: `tjn2@lehigh.edu`

Department of Mathematics, SUNY at Buffalo, Buffalo, NY 14260
E-mail address: `ramac-m@newton.math.buffalo.edu`